

If you have a visual impairment this leaflet can be made available in bigger print or on audiotape. If you require either of these options please contact the Patient Information Centre on 0161 922 5332

#### Language, Interpretation and Patient Support Service (LIPS):

If you require an interpreter to assist your appointment, please ask an appropriate family member to contact our central booking office between **Monday to Friday 8am to 5pm** on **0161 922 6991** to arrange this for you. Further information can be found on the Trust public website <https://www.tamesidehospital.nhs.uk/patients/lips.htm>

#### 語言翻譯及病者支持服務 (LIPS):

如果閣下需要翻譯員在您的預約當日幫助您的話 請找一名合適的家庭成員 **0161 922 6991** 聯絡本中央預約辦事處來您您安排 我們的辦公時間是星期一至星期五 上午 8 時至下午 5 時

#### Językowo Tłumaczeniowa Usługa Pomocy dla Pacjenta (Language, Interpretation and Patient Support Service LIPS):

Jeśli potrzebujesz pomocy tłumacza w trakcie swojej wizyty, proszę poprosić odpowiedniego członka rodziny o skontaktowanie się z Centralnym Biurem Zamówień (Central Booking Office), w celu zorganizowania tłumacza pomiędzy poniedziałkiem a piątkiem w godzinach od 08:00 - 17:00 pod numerem **0161 922 6991**.

لینگویج، انٹریپٹیشن اینڈ پیڈیٹنٹ سپورٹ سروس (Lips)

اگر آپ کو اپنی اپائنٹمنٹ کے لئے مترجم کی مدد کی ضرورت ہو تو براہ مہربانی اپنے خاندان کے کسی موزوں فرد سے کہیں کہ وہ ہمارے سنٹرل بک آفس سے پیر سے جمعہ 8.00 بجے صبح سے 5.00 بجے شام کے دوران 0161 922 6991 پر فون کر کے اس کا بندوبست کریں۔

**Help us to help you** It is important that we keep your records up-to-date. If the information about you is incorrect, we may be unable to contact you should we need to inform you about any changes to your appointment. Therefore, if you change your GP/Dentist, address or telephone number, please contact us as soon as possible. Please provide a mobile number where possible as we operate an appointment reminder service via text message.

Author: Infection Prevention Team©  
Division: Specialist Nursing  
Version No: 4.0  
Ref No: IC022  
Created: August 2011  
Date of last review: June 2020  
Next review date: June 2022



# Tameside and Glossop Integrated Care

## NHS Foundation Trust



## Clostridium Difficile (CDT)

### Discharge from Hospital

This leaflet is intended for patients who have had *Clostridium difficile* infection while in hospital and are now going home. It also contains information for relatives and carers.

If you have been discharged from hospital because you are clinically well but are still on treatment for your *Clostridium difficile* infection, it is important that you follow the instructions given to you by your doctor.

**Things you need to know**

- There is a 20%-30% chance that the *Clostridium difficile* infection and the associated diarrhoea may return (termed a relapse).
- *Clostridium difficile* produces spores (which are structures in this bacterial germ that are resistant to antibiotics) that can survive for several weeks in the environment and on surfaces.
- Spores can be spread by the hands and can lead to infection by entering the body through the mouth.

**It is important for everyone to wash their hands frequently with soap and water to minimize the risk of spreading infection.**

- Your eating pattern does not need to change – just continue with a normal healthy diet. Probiotic drinks may be beneficial.
- Maintain good hand hygiene, by washing your hands with soap and water especially after visiting the toilet and before preparing and eating meals.
- You can return to work if you have been free from diarrhoea for 48 hours. Family members will not be a risk to you.
- Follow strict hand hygiene and practices. If you require antibiotics for any other illness, ensure that your doctor knows that you have had a *Clostridium difficile* infection. As a rule your doctor should only prescribe a short course of antibiotics usually of 3 days duration or less. Show your doctor the green *Clostridium difficile* Alert card

**People in the following groups are more at risk of developing *Clostridium difficile*:**

- Over 65 years old;
- Taking, or who have recently taken, antibiotics or other medication altering the normal bacteria in the gut;
- With a reduced resistance to infection generally;
- Who have spent a long time in a healthcare setting;
- With a serious underlying illness; and
- Who have had bowel surgery

A very small number of people suffer multiple relapses; further treatments must be discussed with your doctor, a further sample for *clostridium difficile* is not usually required.

**If you develop diarrhea at home**

- Contact your GP immediately and ensure that they are aware that you have previously been diagnosed as having *Clostridium difficile* infection.
- Do not take any medicines to stop the diarrhoea (anti-diarrhoeals) while you are suffering from *Clostridium difficile*, when advised to do so by your GP.
- Wash your hands frequently with soap and water.
- Soiled items should ideally be washed at 60°C, or as hot a wash as the fabric can withstand. Take care not to overload the machine so that the water can circulate freely, as this will help to rinse away any germs.
- Remember to wash your hands with soap and water after handling soiled items.
- In the event of heavy faecal soiling clean the excess faeces off soft furnishings with detergent and water and if necessary these can be steamed cleaned. For hard surface contamination a bleach product designed for hard surface disinfection can be used.
- If you use incontinence pads and they are heavily soiled and require disposal. Please contact your GP for advice.
- Hand hygiene with soap and water will break the chain of infection and helps to reduce the spread of infection.

**If you have any questions you want to ask, you can use this space below to remind you.**

Reference: Department of Health (2010). C.Difficile-now you are going home. [www.dh.gov.uk/publications](http://www.dh.gov.uk/publications)